

Article

Synchronous Evolution of an Odor Biosynthesis Pathway and Behavioral Response

Qian Li,¹ Wayne J. Korzan,¹ David M. Ferrero,¹ Rui B. Chang,¹ Dheeraj S. Roy,¹ Mélanie Buchi,² Jamie K. Lemon,¹ Angeldeep W. Kaur,³ Lisa Stowers,³ Markus Fendt,^{2,4} and Stephen D. Liberles^{1,*}

¹Department of Cell Biology, Harvard Medical School, Boston, MA 02115, USA

²Novartis Institutes for BioMedical Research, Neuroscience DA, 4056 Basel, Switzerland

³Department of Cell Biology, The Scripps Research Institute, La Jolla, CA 92037, USA

Summary

Background: Rodents use olfactory cues for species-specific behaviors. For example, mice emit odors to attract mates of the same species, but not competitors of closely related species. This implies rapid evolution of olfactory signaling, although odors and chemosensory receptors involved are unknown.

Results: Here, we identify a mouse chemosignal, trimethylamine, and its olfactory receptor, trace amine-associated receptor 5 (TAAR5), to be involved in species-specific social communication. Abundant (>1,000-fold increased) and sex-dependent trimethylamine production arose de novo along the *Mus* lineage after divergence from *Mus caroli*. The two-step trimethylamine biosynthesis pathway involves synergy between commensal microflora and a sex-dependent liver enzyme, flavin-containing monooxygenase 3 (FMO3), which oxidizes trimethylamine. One key evolutionary alteration in this pathway is the recent acquisition in *Mus* of male-specific *Fmo3* gene repression. Coincident with its evolving biosynthesis, trimethylamine evokes species-specific behaviors, attracting mice, but repelling rats. Attraction to trimethylamine is abolished in TAAR5 knockout mice, and furthermore, attraction to mouse scent is impaired by enzymatic depletion of trimethylamine or TAAR5 knockout.

Conclusions: TAAR5 is an evolutionarily conserved olfactory receptor required for a species-specific behavior. Synchronized changes in odor biosynthesis pathways and odor-evoked behaviors could ensure species-appropriate social interactions.

Introduction

Many animals must effectively distinguish species-specific odors to ensure that mating and other social behaviors are properly directed. In some insect species, changes in pheromone blend compositions are sufficient to drive sexual isolation and speciation [1, 2]. Such evolutionary events require concurrent adaptations in neural systems that control pheromone-evoked behaviors [1].

In the mouse, signaling in the olfactory epithelium (OE) is essential for several odor-driven social behaviors, including

mating, aggression, and nurturing of young [3–6]. Olfactory sensory neurons detect odors and pheromones using a large repertoire of chemosensory receptors, including ~1,100 odorant receptors and 14 trace amine-associated receptors (TAARs) [7]. Whereas receptors for natural mouse odors have been identified in the vomeronasal organ (VNO) [8, 9], TAAR5 is the only known OE receptor activated by a sexually dimorphic mouse odor [10]. TAAR5 detects male mouse odor with exquisite sensitivity, and this response is both sex and age dependent [10].

Pharmacological screens involving hundreds of chemically diverse odorants identified trimethylamine as a high affinity TAAR5 agonist [10]. Trimethylamine is a bacterial metabolite found in some animal odors, and to humans it is a repulsive odor associated with bad breath and spoiled food [11]. Furthermore, the human genetic disease trimethylaminuria is characterized by abnormally high excretion rates of trimethylamine and a resulting body odor that is profoundly aversive [11]. The molecular basis for this disease is mutation of flavin-containing monooxygenase 3 (FMO3) [12], an enzyme that removes endogenous trimethylamine odor by oxidation to form nonvolatile trimethylamine oxide. Trimethylaminuria is a socially debilitating disease, and patients can ameliorate symptoms by reducing dietary intake of choline, a metabolic precursor of trimethylamine, or by taking a regimen of antibiotics [11]. In comparison with humans, mice normally release >1,000-fold more urinary trimethylamine, and they do so in a sex-dependent manner. Furthermore, TAAR5 sensitively detects mouse urine, but does not detect human urine at any concentration examined [10]. This suggests key evolutionary differences in the trimethylamine biosynthesis pathways of mouse and human.

Here, we identify specific evolutionary changes in this pathway that occurred along the *Mus* lineage. In mouse, the principal trimethylamine oxidase, FMO3, is expressed at >1,000-fold higher levels in female liver than male liver. In contrast, closely related rodent species produce low or undetectable levels of trimethylamine, and neither trimethylamine production nor *Fmo3* gene expression is sex dependent. Sexually dimorphic trimethylamine production is most distantly observed in *Mus spretus*, a species that recently diverged from *Mus musculus*. The recent and de novo evolution of sex-dependent *Fmo3* gene control in mice provides a molecular basis for changes in scent odors between highly related mammalian species.

Furthermore, we provide evidence for coincident evolution of trimethylamine biosynthesis pathways and trimethylamine-evoked behavioral responses in rodents. Several TAAR agonists are aversive chemicals, such as 2-phenylethylamine, a carnivore odor that repels rodents [13, 14]. Trimethylamine is also aversive to humans, but behavioral responses of mice to this chemical have not been characterized. Because trimethylamine is abundant in mouse urine, a richly attractive odor source for mice, we reasoned that trimethylamine might evoke different behavioral responses in mice. Here, we find that the valence of mouse behavioral responses to trimethylamine is highly concentration dependent, with attraction occurring at physiological concentrations found in urine.

*Present address: Institute for Pharmacology and Toxicology, University of Magdeburg, 39120 Magdeburg, Germany

*Correspondence: stephen_liberles@hms.harvard.edu




Figure 1. Species- and Sex-Dependent Production of the TAAR5 Activator Trimethylamine

(A) Variations in production of the TAAR5 activator across mammalian species. HEK293 cells were transfected with plasmids encoding CRE-SEAP and TAAR5 (black bars) or CRE-SEAP alone (white bars), incubated with serial dilutions of urine from the six species indicated, and assayed for SEAP activity (triplicates \pm SD). Rodent donors were males, and human donors were nonidentifiable males and females.

Knockout mice lacking TAAR5 do not display this attraction behavior, and rats, who retain a functional TAAR5, instead avoid trimethylamine at the same concentration. Enzymatic depletion of trimethylamine or disruption of TAAR5 by gene knockout decreases scent attraction in mice. Thus, we demonstrate that an odor response requires a particular OE receptor, and that this response is malleable across species. Synchronized changes in odor biosynthesis pathways and odor-evoked behaviors could ensure species-appropriate social interactions.

Results

TAAR5 Detects a Species-Dependent Odor

Receptors in the olfactory epithelium that discriminate odors of closely related species are unknown. In the course of characterizing TAAR activators across species, we observed robust activation of TAAR5 by mouse odors, but not by odors of related rodents.

TAAR5 responses were measured using an established reporter gene assay [10, 14]. HEK293 cells were cotransfected with plasmids containing TAAR5 and a cAMP-dependent reporter gene (CRE-SEAP), incubated with titrations of test specimens, and assayed for reporter activity. Consistent with our previous findings [10], TAAR5 exhibited robust and sensitive responses to mouse urine (Figure 1A) that were sex dependent (Figure 1B). Male mouse urine evoked threshold and half-maximal (EC_{50}) TAAR5 responses when diluted 100,000-fold and 30,000-fold, respectively.

We examined TAAR5 responses to urine from five other mammalian species: rat, gerbil, hamster, guinea pig, and human. In contrast to the sensitive responses observed to mouse urine, TAAR5 did not detect rat, guinea pig, or human specimens at any dilution tested (up to 100-fold; see Figure S1 available online). Hamster and gerbil urine elicited threshold responses at 300-fold and 1,000-fold dilutions, respectively, and thus contained significantly lower levels of the TAAR5 activator than mouse urine. Furthermore, TAAR5 responses to urine from either male or female rats were not observed. Variations in production of the TAAR5 activator were thus greater between species (>1,000-fold between mouse and rat) than between mouse sexes (~10- to 30-fold). Because multiple rodents do not similarly produce the TAAR5 agonist, a parsimonious interpretation would be that its abundant and sex-dependent production in mouse is a recent gain-of-function adaptation.

Identification of Species- and Sex-Dependent Mouse Odors by NMR Spectroscopy

To probe how TAAR5 distinguishes mouse and rat urine, as well as male and female mouse urine, we characterized abundant species-specific and sex-specific urinary odors by nuclear magnetic resonance (NMR) spectroscopy (Figure 1C).

(B) Assays were performed as above, except donors were male and female mice (triplicates \pm SE).

(C) A region (2.8 to 3.6 ppm) of the NMR spectra of male and female mouse urine (C57BL/6), trimethylamine, and trimethylamine oxide. Chemical assignments were based on published values [15] or analysis of spiked specimens.

(D) qPCR analysis of *Fmo3* gene expression in liver cDNA prepared from mice of sexes and ages indicated (triplicates \pm SE). Copy numbers were calculated in cDNA derived from 10 ng of liver RNA by comparison with PCR reactions involving titrations of an *Fmo3*-containing plasmid.

The largest peak in the NMR spectrum of male mouse urine corresponded to trimethylamine (5 mM), and production of trimethylamine was sex dependent, with ~20-fold enrichment in male urine compared to female urine. This is consistent with data indicating TAAR5 to be a trimethylamine-activated olfactory receptor that discriminates male and female mouse odors [10]. The abundance of trimethylamine is sufficient to explain the striking sensitivity of TAAR5 for male mouse urine. Furthermore, trimethylamine was not detected in male rat urine by NMR spectroscopy (Figure S1), consistent with the inability of rat specimens to activate TAAR5 (Figure 1A).

Comparative NMR spectroscopy of male and female mouse urine also identified a second related chemical produced with opposing sex dependence. Based on analysis of spiked specimens, this female-enriched chemical was trimethylamine oxide, a chemical in the same biosynthetic pathway as trimethylamine [16]. The levels of other major metabolites, including creatine and taurine, were similar between males and females. The reciprocal production of trimethylamine and trimethylamine oxide by males and females suggested trimethylamine oxidation to be a sex-dependent chemical reaction in mice.

Sex-Specific Expression of Trimethylamine Oxidase in the Mouse

FMO3 is the principal trimethylamine oxidase, as mutation of the *Fmo3* gene in humans causes trimethylaminuria [12]. We asked whether differences in FMO3 expression between male and female mice could explain sex-dependent trimethylamine production. It was previously reported that FMO3 expression in mouse liver, its primary site of expression, is regulated by a testosterone-dependent mechanism, although a function for this regulation was not clear [17].

Here, we performed a quantitative analysis of *Fmo3* gene expression in liver cDNA derived from male and female mice of different ages (Figure 1D). We observed a striking >1,000-fold elevation of *Fmo3* gene expression in adult females compared to adult males, which was consistent with previous results [17]. Male-specific silencing of *Fmo3* gene expression is discernible between 4 and 5 weeks of age, matching the time course of trimethylamine appearance in male urine [10]. For comparison, expression of *Gapdh* was similar across all specimens (Figure S1). Expression levels of *Fmo3* in liver are thus inversely correlated with levels of trimethylamine in urine of male, female, and prepubescent mice.

A Two-Step Trimethylamine Biosynthesis Pathway in Mouse Involves Microbial Metabolism and FMO3 Oxidation

The trimethylamine biosynthesis pathway is well studied in humans [18] but exhibits several key differences in mice, where urinary trimethylamine release is sex dependent and elevated >1,000-fold. Trimethylamine is initially derived by bacterial metabolism of dietary choline [19], and trimethylaminuria is ameliorated by decreasing choline intake or by antibiotic treatment [11]. Here, we asked if the normal production of trimethylamine in mouse urine is similarly controlled. Urine was collected from mice fed a choline/methionine-free diet or treated with the antibiotic neomycin sulfate (2 mM in drinking water) for 1 week and verified to have similar concentrations of several common metabolites by NMR spectroscopy (see below).

Specimens were then analyzed for the ability to activate TAAR5 using the reporter gene assay (Figures 2A and 2B). As described above, TAAR5 exhibited robust and sensitive responses to urine of mice fed a standard diet. In contrast,

TAAR5 had ~10-fold reduced sensitivity for the urine of sex-matched mice fed a choline/methionine-free diet or placed on a regimen of antibiotics for 1 week (Figures 2A and 2B).

Next, we performed NMR spectroscopy on urine from diet-altered and normal-fed mice to determine if shifts in TAAR5 activation curves could be explained by reductions in trimethylamine levels (Figures 2C and 2D). Diet alteration did not impact urinary levels of taurine and creatine but reduced levels of both trimethylamine and trimethylamine oxide. Trimethylamine levels were 5 mM in urine from normal-fed males and 400 µM in diet-altered males, a ~12-fold difference that matched observed changes in TAAR5 activation curves (Figure 2A). Antibiotic treatment resulted in a similar reduction in urinary trimethylamine (Figure 2E). The antibiotic treatment and diet modification eliminated most urinary trimethylamine (~90%), but not all. Residual trimethylamine in mouse urine could be derived from metabolism of other nitrogen-containing nutrients or from the endogenous reservoir of phosphatidylcholine lipid. Nevertheless, these data indicate that a major step in the trimethylamine biosynthesis pathway of mice involves the metabolism of dietary choline by endogenous microbes.

Together, these data are consistent with a two-step biosynthetic pathway for trimethylamine in mice: first, trimethylamine is derived largely via metabolism of dietary choline by gut flora, and second, trimethylamine odor is reduced in females, but not males, by a sex-dependent oxidation reaction involving FMO3 (Figure 2F).

Evolution of a Sexually Dimorphic Biosynthesis Pathway for a Mouse Odor

We next sought to understand the evolutionary origins of abundant and sex-dependent trimethylamine biosynthesis along the *Mus* lineage (Figure 3A). We obtained urine from five highly related *Mus* species: *Mus musculus*, *Mus caroli*, *Mus spicilegus*, *Mus spretus*, and *Mus domesticus*. Specimens were collected from age- and sex-matched animals housed under similar environmental conditions and fed standard diets with identical levels of choline. Fresh collections were used to prevent odor evaporation or bacterial growth, and samples were verified to have similar levels of creatinine (Figure S2). TAAR5 responses were then measured to serial dilutions of each specimen over a large concentration range, using our cellular reporter gene assay (Figure 3B).

Mus caroli specimens, like *Rattus norvegicus* specimens, did not activate TAAR5 at any concentration tested (up to 100-fold, Figure S1). Based on these data, *Mus musculus* males emit >1,000-fold higher levels of trimethylamine than either of these species. In contrast, TAAR5 did detect urine from *Mus spretus* and *Mus spicilegus* but did so with reduced sensitivity. These species produce lower amounts of trimethylamine than *Mus musculus* but higher amounts than *Rattus norvegicus* (Figure S2), and could represent evolutionary intermediates in this biosynthesis pathway. Sexually dimorphic trimethylamine production is most distantly observed in *Mus spretus* and maintained in both *Mus domesticus* and *Mus musculus*. Lower levels of sexual dimorphism are observed in *Mus spicilegus*; males and females have slight differences in the production ratio of trimethylamine to creatinine (Figure S2), but the extent of sexual dimorphism is reduced in this species. Together, these data indicate that abundant and sex-dependent production of trimethylamine arose recently along the *Mus* lineage and likely within the Eurasian clade of mice.




Figure 2. Trimethylamine Biosynthesis in Mouse Involves Microbial Metabolism and a Sex-Dependent Oxidation Reaction

Urine was collected from male animals provided with normal chow, choline/methionine-free diet, or antibiotics for 1 week. Trimethylamine levels in these specimens were determined by the TAAR5-mediated reporter gene assay (A and B; triplicates \pm SD) and by NMR spectroscopy (C–E). These data are consistent with a two-step pathway regulating trimethylamine production in mouse (F).

trimethylamine production. *Fmo3* gene expression was sexually dimorphic, with a male to female expression ratio of $>1,000$ in four strains of *Mus musculus* (Figure S2) and >100 in *Mus domesticus*, a closely related subspecies. Sexually dimorphic *Fmo3* gene expression is most distantly observed in *Mus spretus*, and it is not similarly observed in *Mus caroli* or *Rattus norvegicus*. These data indicate that *Fmo3* gene repression in males is a recent alteration that occurs in those *Mus* species with sexually dimorphic trimethylamine production.

In addition, we found that *Fmo3* expression in the kidney also varied between related rodent species (Figure 3D). Interestingly, *Fmo3* gene expression in kidney was not sex specific in any species, and it was ~ 100 -fold lower in *Mus musculus* than *Rattus norvegicus*. This second evolutionary change in *Fmo3* expression likely explains why trimethylamine levels differ between female mice and female rats, despite similar levels of liver *Fmo3*.

Trimethylamine Attracts Mice at Concentrations Normally Found in Mouse Urine

We present evidence that trimethylamine is abundant in mouse urine, and


that multiple genetic changes have occurred in the trimethylamine biosynthesis pathway during *Mus* evolution. These data suggest that trimethylamine could be a species-specific chemosignal for mice, although associated behavioral responses have not been investigated. We reasoned that trimethylamine could be either attractive to mice, like mouse urine that contains chemicals important for social behaviors, or aversive to mice, like trimethylamine to humans or other TAAR agonists, 2-phenylethylamine and isoamylamine, to rodents [11, 13, 21].

We measured the valence of trimethylamine-evoked behavioral responses in mice using a two-choice compartment assay based on previously reported paradigms [13, 21]. Briefly, mice were introduced into a two-compartment test arena (Figure 4A), with one compartment containing test stimuli ($400 \mu\text{l}$) in a volatile odor applicator. The two compartments were separated by a curtain, and odor plumes were controlled to prevent mixing across compartments. Time

FMO3 Expression Is an Evolutionary Node in the Trimethylamine Biosynthesis Pathway

FMO3 is a key regulator of trimethylamine production [12], so we asked whether FMO3 expression varied across *Mus* species in accordance with changes in trimethylamine production. We prepared liver and kidney cDNA from male and female animals of several species, including *Rattus norvegicus*, *Mus caroli*, *Mus spilelegus*, *Mus spretus*, *Mus domesticus*, and four strains of *Mus musculus*. Levels of *Gapdh* mRNA were similar in all of these specimens (Figure S2). We then determined *Fmo3* gene expression levels by qPCR, using primers that recognized a highly conserved region of *Fmo3* coding sequence.

We found that *Fmo3* gene expression levels in liver were sexually dimorphic in some, but not all, *Mus* species (Figure 3C). Differences in *Fmo3* gene expression between species were primarily due to differences in male-specific *Fmo3* gene silencing. Moreover, there was a strong correlation between sex-dependent *Fmo3* gene repression and sex-dependent






Figure 3. Evolution of Sex-Dependent Trimethylamine Biosynthesis and *Fmo3* Gene Repression along the *Mus* Lineage

(A) The phylogenetic relationship of *Mus* species, with *Rattus norvegicus* as an outgroup. The figure is adapted from published data [20]. *Mus musculus* subspecies are shown in the gray box, and bracketed letters indicate species providing specimens for subsequent panels.

(B) Urine from males (gray bars) and females (black bars) of species indicated were collected and analyzed using the TAAR5 reporter gene assay.

(C and D) *Fmo3* gene expression in liver and kidney of various rodents. Liver and kidney cDNA was prepared from species indicated, as coded by bracketed letters in the phylogenetic tree (A), and analyzed for *Fmo3* gene expression by qPCR analysis (triplicates \pm SE). *Fmo3* expression levels in male animals (gray bars) and female animals (black bars) were compared. The y axis indicates copy number in cDNA derived from 10 ng RNA and was calculated by comparison with PCR reactions involving titrations of an *Fmo3*-containing plasmid.




Figure 4. Biphasic Valence of Mouse Behavioral Responses to Trimethylamine

(A) A cartoon depiction of the mouse behavioral assay.

(B) Side preferences of mice in a two-compartment arena containing test odors and water were measured as differential occupancy. Increases in test-odor-side occupancy were scored as attraction, and decreases were scored as aversion (* $p < 0.05$, ** $p < 0.01$, \pm SE). All experiments involved equal numbers of males and females, except as noted in [Supplemental Experimental Procedures](#).

spent on each side of the arena was recorded during a 2 min test period, and differences in compartment occupancy were scored as attraction or aversion. This experimental design was validated with control odors (Figure 4B); opposite sex urine, a powerful attractant for mice, increased test compartment occupancy (+43%), whereas a fox-derived predator odor, 2,5-dihydro-2,4,5-trimethylthiazole (TMT) [22], decreased test compartment occupancy (−45%). Furthermore, water and eugenol, a novel odor, neither increased nor decreased test compartment occupancy significantly in this paradigm (water −1.5%; 10 mM eugenol +2.3%, 4.2 M eugenol +2.7%).

Trimethylamine evoked biphasic and concentration-dependent behavioral responses in mice, with aversion occurring to high trimethylamine concentrations and attraction occurring to lower, physiologically relevant trimethylamine concentrations (Figure 4B). Decreased test compartment occupancy was observed to trimethylamine at 4.2 M (−28%, $n = 20$, $p < 0.001$) and 1 M (−19%, $n = 16$, $p = 0.03$), but increased test compartment occupancy was observed to trimethylamine at 100 mM (+26.4%, $n = 20$, $p < 0.001$), 10 mM (+20.7%, $n = 32$, $p = 0.001$), and 1 mM (+11.7%, $n = 20$, $p = 0.07$). At all concentrations, trimethylamine responses were indistinguishable in male and female animals (Figure S3). Thus, trimethylamine, in the absence of other urinary cues, attracts mice at concentrations normally found in mouse urine. We also examined a role for trimethylamine in other behavioral paradigms, but observed partial or no effect in the resident-intruder

aggression paradigm and various odor discrimination tasks (Figure S3).

Trimethylamine Is Aversive to Rats

Although trimethylamine is attractive to mice, it is highly aversive to humans. Furthermore, trimethylamine production is elevated >1,000-fold in mice than in rats, so we reasoned that rats and mice could display distinct behavioral responses to this chemical. We previously published an open-field quadrant assay to measure the valence of odor responses in rats, and we used this assay to demonstrate avoidance of predator odors, including the TAAR4 agonist 2-phenylethylamine [13, 23]. As with the mouse paradigm, we observed similar responses to control stimuli, including neutral responses to water (24% quadrant occupancy or qo) and attraction responses to opposite sex urine (35% qo).

Unlike mice, rats were not attracted to any concentration of trimethylamine tested (Figures 5A and 5B). Rats exhibited decreases in odor-quadrant occupancy when trimethylamine was used as a stimulus at 4.2 M (1.7% qo), 1 M (4.3% qo), 100 mM (11.2% qo), and 10 mM (14.2% qo). We next investigated behavioral responses of rats to mouse urine, an abundant source of trimethylamine. Rats are predators of mice, so we reasoned that mouse urine, as prey odor, could contain either attractive kairomones intercepted by rats or aversive allomones released by mice for defensive behavior. Using the open-field quadrant assay, we observed that male mouse urine was aversive to rats and that female urine became aversive when male levels of trimethylamine were added (Figure 5C). These data suggest a function for trimethylamine as an aversive allomone and demonstrate that trimethylamine evokes responses of different valence in mice and rats.

We next asked whether pharmacological differences in TAAR5 orthologs from mouse and rat could explain variations in the valence of trimethylamine responses. However, using the reporter gene assay (Figure 5D), we observed that both rat and mouse TAAR5 detected trimethylamine with similar sensitivity (mouse EC₅₀ = 300 nM; rat EC₅₀ = 1 μM). Furthermore, single-neuron imaging and post hoc expression analysis determined that mouse TAAR5, in the context of the olfactory sensory neuron, also detected trimethylamine at 300 nM, but not isoamylamine or 2-phenylethylamine, agonists for TAAR3 and TAAR4 (Figure S4). Trimethylamine thresholds that evoke behavioral responses are understandably different from those that activate TAAR5 in heterologous cells, because our behavioral assays require investigation of a trimethylamine odor source from a distance. It is possible that these species-particular odor responses involve differences in olfactory circuits downstream of receptor activation. Alternatively, trimethylamine may activate different repertoires of receptors in mouse and rat, and receptors other than TAAR5 could mediate either mouse attraction or rat aversion.

TAAR5 Is Required for Mouse Attraction to Trimethylamine

Although several TAAR agonists are aversive odors, trimethylamine evokes a more complex and distinct response in mice. We next asked if TAAR5 was required for the avoidance response to high levels of trimethylamine or the attraction response to physiological levels of trimethylamine. We obtained TAAR5 knockout mice (*Taar5*^{−/−}) from the Knockout Mouse Project (KOMP) repository [24], and *Taar5* expression was undetectable in olfactory epithelium derived from homozygous animals by *in situ* hybridization (Figure 6A). Next, we measured the valence of behavioral responses to




Figure 5. Rats Avoid a Trimethylamine Odor Source

(A) Motion tracks (red) of individual male rats behaving in the open-field paradigm. White circles indicate the location of odor sources (water, 4.2 M trimethylamine).

(B and C) Occupancy of quadrants, defined as 25% of the arena, containing odors indicated ($n = 9-12$, * $p < 0.05$, ** $p < 0.01$, \pm SE).

(D) Dose-dependent trimethylamine responses of HEK293 cells transfected with reporter gene alone (white squares), reporter gene and mouse TAAR5 (black squares), or reporter gene and rat TAAR5 (gray squares) were measured using the cellular reporter gene assay (triplicates \pm SE).

trimethylamine in *Taar5*^{-/-} animals, using the odor compartment assay (Figure 6B). We observed that *Taar5*^{-/-} animals displayed decreased attraction to trimethylamine at all concentrations tested. At high concentrations (4.2 M), avoidance behavior was more pronounced because occupancy of *Taar5*^{-/-} animals in the odor compartment was lower (~61.5%, $n = 12$) than wild-type animals (~28%, $n = 20$). Furthermore, at concentrations that attracted wild-type mice, trimethylamine evoked neither attraction nor aversion responses in *Taar5*^{-/-} animals, including at 100 mM

(~11.1%, $n = 8$) and 10 mM (~2.2%, $n = 14$). These responses were statistically indistinguishable from neutral responses to control odors. Thus, TAAR5 is a single mouse OE receptor required for attraction to trimethylamine.

Trimethylamine and TAAR5 Function in Scent Attraction

Next, we asked if the ligand-receptor pair of trimethylamine and TAAR5 is required for attraction to mouse urine. We developed a strategy to deplete trimethylamine from mouse urine based on its biosynthesis pathway. Male mouse urine was incubated with or without FMO3-containing microsomes to create “TMA-depleted male urine” and “mock-depleted male urine.” Trimethylamine levels were decreased >300-fold in TMA-depleted male urine, as determined by an inability to activate TAAR5 across a range of concentrations (Figure 6C). Although FMO3 has high substrate specificity for trimethylamine [25], we reasoned that FMO3 could potentially oxidize other urinary chemicals under these reaction conditions. To control for this, we created an additional test specimen (“TMA-respiked male urine”), in which trimethylamine was restored to normal levels following removal of FMO3 microsomes by centrifugation.

Next, we examined if TMA-depleted, mock-depleted, and TMA-respiked male urine (40 μ l) were similarly attractive to mice using the two-choice compartment assay (Figure 6D). Mock-depleted urine was a highly attractive stimulus (+37.8%, $n = 18$), similar to what we observed for intact urine (Figure 4). In contrast, mice exhibited decreased attraction toward TMA-depleted male urine (+17.8%, $n = 20$), and this decrease persisted over several urine doses tested (Figure S5). Furthermore, attraction was restored to TMA-respiked male urine (+30.8%, $n = 20$). Simultaneous presentation of mock-depleted and TMA-depleted urine in the two-choice arena also indicated a preference for mock-depleted urine (Figure S5). Finally, *Taar5*^{-/-} animals showed reduced attraction to mock-depleted male urine (+17.1%, $n = 10$) that was not seen in sex- and age-matched wild-type controls (Figure 6E). These data indicate a role for trimethylamine and TAAR5 in attraction to mouse scent, and they also indicate that there are other attractive olfactory cues in male mouse urine. Thus, mouse scent is an attractive odor blend containing multiple chemicals, among them trimethylamine, a species-dependent odor with a rapidly evolving biosynthesis pathway.

Discussion

Animals have evolved elaborate and diverse sex traits important for species-specific behaviors. Rodents use olfactory cues as such traits, and both odor production pathways and odor-evoked behavioral responses are under strong evolutionary pressure for diversification between related species. We previously discovered an olfactory receptor, TAAR5, that detects an abundant mouse odor, trimethylamine [10]. Here, we show that both trimethylamine biosynthesis pathways and trimethylamine-evoked behavioral responses vary considerably between related rodents.

Stepwise Evolution of the Trimethylamine Biosynthesis Pathway

Mechanisms by which volatile scent odor biosynthesis pathways have evolved in rodents to include sex-specific and species-specific chemical reaction steps are unknown. In invertebrates, alterations to odor biosynthesis pathways that include changes in the function or expression pattern of




Figure 6. TAAR5 and Trimethylamine Are Required for Mouse Scent Attraction

(A) Expression of *Taar5* and *lacZ* in olfactory sensory neurons of *Taar5^{-/-}*, *Taar5^{+/-}*, and *Taar5^{+/+}* mice visualized by two-color *in situ* hybridization. Scale bar = 50 μ m.

(B) Responses of *Taar5^{+/+}* and *Taar5^{-/-}* mice to trimethylamine were measured, using the two-choice odor-compartment assay (mean \pm SE).

specific biosynthetic enzymes can drive the evolution of new species [1, 2, 26]. In European corn borer moths, altered substrate recognition of a fatty-acyl reductase changes the predominant lipid acetate isomer in a pheromone blend [2]. This change in enzyme function can cause reproductive isolation, a first step in speciation. In *Drosophila*, rapid changes in the expression patterns of desaturases correlate with species-particular differences in the production of long-chain cuticular hydrocarbons that function as sex pheromones [26].

Here, we identify FMO3-catalyzed oxidation to be an evolutionary node in the trimethylamine biosynthesis pathway of rodents. In humans, rats, and other species, FMO3 normally clears trimethylamine odor by conversion to nonvolatile trimethylamine oxide [12, 18]. Mice have evolved specific mechanisms for decreasing the efficiency of this oxidation reaction and enhancing trimethylamine release >1,000-fold, compared to rats. The trimethylamine pathway has undergone stepwise alteration in the *Mus* genus, with *Mus spicilegus* and *Mus spreitus* representing intermediates. Furthermore, abundant and sex-specific trimethylamine production arose de novo after divergence from *Mus caroli*.

How could sex-dependent FMO3 expression arise along the *Mus* lineage? FMO3 is highly expressed in the liver, and sex-dependent liver gene expression predominantly occurs through a growth hormone-mediated mechanism [27]. In mice, androgen signaling modulates the release dynamics of growth hormone from the pituitary; males release growth hormone in rhythmic bursts, whereas females release growth hormone at lower, constant levels. Pulsatile growth hormone release in males enables activation and deactivation cycles of growth hormone-mediated signaling cascades not observed in females. One critical growth hormone-activated transcription factor in liver is Stat5b, which is required for sex-dependent induction of some male-enriched proteins, such as major urinary proteins (MUPs) that function as mouse pheromones, and repression of other female-enriched proteins [28]. Mechanisms by which Stat5b activates some genes but silences others are not understood [27] and could involve transcriptional coactivators and corepressors that are not yet identified. Sequence analysis of the *Fmo3* promoter across species (Supplemental Experimental Procedures) identified several regions upstream of the *Fmo3* transcription start site that encoded transcription-factor binding sites and varied across rodents in accordance with sex-dependent trimethylamine production and *Fmo3* gene repression. One site that arose de novo is a consensus binding site for Stat5b; other mutations created binding sites for transcription factors with no known and direct function in sex-dependent liver gene expression. The identification of novel transcription-factor binding sites in the *Fmo3* gene promoter that arose concurrently with sex-dependent *Fmo3* gene repression is potentially interesting for future study because Stat5b corepressors that mediate male-specific gene silencing in liver are unknown.

Experiments involved equal numbers of males and females, except as noted in Supplemental Experimental Procedures.

(C) Male urine was depleted of trimethylamine by incubation with FMO3-containing microsomes. Effective depletion was shown by an inability of TMA-depleted male urine to activate TAAR5 in HEK293 cells at dilutions indicated (triplicates \pm SE).

(D) Behavioral responses of mice to stimuli indicated (40 μ l), using the two-choice compartment assay (mean \pm SE).

(E) Behavioral responses of female mice ($n = 8\text{--}10$, \pm SE) of genotypes indicated to mock-depleted male urine (40 μ l) (* $p < 0.05$, ** $p < 0.01$).

Bacteria and Scent Odors

These studies highlight an important role for endogenous microflora in generating odor cues that influence mouse scent composition. In humans, bacterial metabolites influence the odor quality of axillary sweat and other secretions [29]. In rodents, bacterial and dietary metabolites could provide a mechanism for distinguishing urinary odors of genetically identical animals [30, 31] and could be dynamic in wild mice, varying with ecological niche. In laboratory mice, under homogenous conditions, levels of trimethylamine production were not highly variable between adult male animals.

In contrast to odors generated by endogenous bacteria, odors generated by invasive bacteria can provide repulsive signals that indicate infection or illness [32]. Formyl peptide receptor-related sequences function as sensory receptors in the vomeronasal organ, and they are related to immune system receptors that recognize formylated peptides released at sites of tissue damage and bacterial infection [33, 34]. Furthermore, solitary chemosensory cells in the mouse respiratory epithelium contain bitter taste receptors that detect acyl-homoserine lactones involved in bacterial quorum sensing [35]. Thus, mice have evolved numerous chemosensory mechanisms for recognizing endogenous, invasive, or environmental bacteria.

Mouse Attraction to a Trimethylamine Odor Source Requires TAAR5

Trimethylamine is abundant in mouse urine; highly volatile; produced with sex, age, and species dependence; and activates a mouse olfactory receptor with high affinity. We also find that trimethylamine (10 mM) is an attractive cue to mice, and mouse urine depleted of trimethylamine evokes decreased attraction responses. Trimethylamine is one of several attractive components of mouse urine, consistent with previous imaging and behavioral studies indicating that mouse odor recognition involves detection of a blend of constituents [36, 37]. Other attractive components in mouse urine include volatiles such as (methylthio)-methanethiol, 2-sec-butyl-dihydrothiazole, dehydro-exo-brevicomin, and farnesenes, as well as a nonvolatile MUP termed darcin [36–39].

In our paradigm, male and female mice display similar levels of attraction to trimethylamine (Figure S3). It is possible that trimethylamine evokes a sex-specific behavioral response that we have not identified, and in future studies, it would be interesting to examine a role for trimethylamine and TAAR5 in influencing sexual receptivity of females. Alternatively, trimethylamine could provide a general attraction signal, with selective pressure for enriched production by males arising because scent marking is a stereotyped territorial behavior typically displayed by male animals.

Mouse attraction responses to trimethylamine require a particular OE receptor, TAAR5. There are only a few examples where alterations of individual olfactory receptors in mammals change odor-evoked behaviors. In mice, genetic deletion of a cluster of VNO receptors influences some social behaviors [40], and deletion of a single VNO receptor effects a specific mating response to the male lacrimal pheromone ESP1 [8]. Furthermore, polymorphisms in a human odorant receptor for androstenone account for changes in human odor perception [41]. Here, we show that knockout of a specific OE receptor in mice alters behavioral responses to the volatile mouse odor trimethylamine.

Trimethylamine Evokes Different Behavioral Responses in Mice and Rats

Rat and mouse TAAR5 similarly detect trimethylamine, yet TAAR5-mediated attraction behavior is only observed in mice. Because rats avoid trimethylamine, it is possible that trimethylamine functions in a defensive capacity as an allomone that reduces attraction of predators or competing species to mouse scents. Consistent with this notion, rats avoid male mouse urine, an abundant source of trimethylamine.

How do the mouse and rat olfactory systems mediate distinct trimethylamine responses? Because several TAAR ligands are aversive to rodents, and trimethylamine is aversive to both humans and rats, a parsimonious interpretation would be that TAAR5-mediated attraction is a recent adaptation in mouse. Adaptive changes in TAAR5-activated olfactory circuits could occur downstream of receptor activation and involve either developmental alterations to hard-wired neural circuits or learning. A role for learning in adaptive behavioral responses simplifies how a species-specific odor response might evolve, because a single change in an odor biosynthesis pathway could be sufficient to drive rapid evolution of olfactory communication. Alternatively, TAAR5 could couple to similar circuits in all species, and rats and humans might have an additional trimethylamine-activated olfactory receptor not found in mice that mediates dominant aversion responses. Consistent with the existence of additional mechanisms for trimethylamine sensing, *Taar5*^{-/-} animals do retain avoidance of high trimethylamine concentrations.

Here, we identify an olfactory receptor that is both evolutionarily conserved and required for a species-specific attraction response. The ligand for this olfactory receptor is the product of a rapidly evolving biosynthesis pathway. *Fmo3* gene control provides a key evolutionary node in this pathway and a molecular basis for changes in odor production between highly related mammalian species. Synchronous evolution of biosynthesis pathways and evoked responses could help ensure that social behaviors are properly targeted to animals of the same species.

Experimental Procedures

A description of reagents and methods for TAAR5 functional assays, qPCR, NMR spectroscopy, mouse behavior, rat behavior, odor depletion, single-neuron calcium imaging, single-neuron expression analysis, *in situ* hybridization, specimen collection, animal husbandry, and statistical analysis are detailed in the *Supplemental Information*.

Supplemental Information

Supplemental Information includes five figures and Supplemental Experimental Procedures and can be found with this article online at <http://dx.doi.org/10.1016/j.cub.2012.10.047>.

Acknowledgments

We thank Sandeep Robert Datta for helpful advice and use of his mouse behavioral suite. We also thank Tom Rapoport, Mark Albers, and Sandeep Robert Datta for valuable comments on the manuscript. We thank the Nikon Imaging Center and NMR Facility at Harvard Medical School. This work was supported by a grant from the National Institute on Deafness and Other Communicative Disorders (S.D.L., award number R01DC010155). D.M.F. is supported by a Boehringer Ingelheim Fonds PhD Fellowship.

Received: July 10, 2012

Revised: October 12, 2012

Accepted: October 25, 2012

Published: November 21, 2012

References

1. Smadja, C., and Butlin, R.K. (2009). On the scent of speciation: the chemosensory system and its role in premating isolation. *Heredity (Edinb)* **102**, 77–97.
2. Lassance, J.M., Groot, A.T., Liénard, M.A., Antony, B., Borgwardt, C., Andersson, F., Hedenström, E., Heckel, D.G., and Löfstedt, C. (2010). Allelic variation in a fatty-acyl reductase gene causes divergence in moth sex pheromones. *Nature* **466**, 486–489.
3. Belluscio, L., Gold, G.H., Nemes, A., and Axel, R. (1998). Mice deficient in G(olf) are anosmic. *Neuron* **20**, 69–81.
4. Mandiyan, V.S., Coats, J.K., and Shah, N.M. (2005). Deficits in sexual and aggressive behaviors in Cnga2 mutant mice. *Nat. Neurosci.* **8**, 1660–1662.
5. Wang, Z., Balet Sindreu, C., Li, V., Nudelman, A., Chan, G.C., and Storm, D.R. (2006). Pheromone detection in male mice depends on signaling through the type 3 adenyl cyclase in the main olfactory epithelium. *J. Neurosci.* **26**, 7375–7379.
6. Wang, Z., and Storm, D.R. (2011). Maternal behavior is impaired in female mice lacking type 3 adenyl cyclase. *Neuropharmacology* **56**, 772–781.
7. Nei, M., Niimura, Y., and Nozawa, M. (2008). The evolution of animal chemosensory receptor gene repertoires: roles of chance and necessity. *Nat. Rev. Genet.* **9**, 951–963.
8. Haga, S., Hattori, T., Sato, T., Sato, K., Matsuda, S., Kobayakawa, R., Sakano, H., Yoshihara, Y., Kikusui, T., and Touhara, K. (2010). The male mouse pheromone ESP1 enhances female sexual receptive behaviour through a specific vomeronasal receptor. *Nature* **466**, 118–122.
9. Isogai, Y., Si, S., Pont-Lezica, L., Tan, T., Kapoor, V., Murthy, V.N., and Dulac, C. (2011). Molecular organization of vomeronasal chemoreception. *Nature* **478**, 241–245.
10. Liberles, S.D., and Buck, L.B. (2006). A second class of chemosensory receptors in the olfactory epithelium. *Nature* **442**, 645–650.
11. Mitchell, S.C., and Smith, R.L. (2001). Trimethylaminuria: the fish malodor syndrome. *Drug Metab. Dispos.* **29**, 517–521.
12. Dolphin, C.T., Janmohamed, A., Smith, R.L., Shephard, E.A., and Phillips, I.R. (1997). Missense mutation in flavin-containing mono-oxygenase 3 gene, FMO3, underlies fish-odour syndrome. *Nat. Genet.* **17**, 491–494.
13. Ferrero, D.M., Lemon, J.K., Fluegge, D., Pashkovski, S.L., Korzan, W.J., Datta, S.R., Spehr, M., Fendt, M., and Liberles, S.D. (2011). Detection and avoidance of a carnivore odor by prey. *Proc. Natl. Acad. Sci. USA* **108**, 11235–11240.
14. Ferrero, D.M., Wacker, D., Roque, M.A., Baldwin, M.W., Stevens, R.C., and Liberles, S.D. (2012). Agonists for 13 trace amine-associated receptors provide insight into the molecular basis of odor selectivity. *ACS Chem. Biol.* **7**, 1184–1189.
15. Gavaghan McKee, C.L., Wilson, I.D., and Nicholson, J.K. (2006). Metabolic phenotyping of nude and normal (Alpk:AfpCD, C57BL10J) mice. *J. Proteome Res.* **5**, 378–384.
16. Wang, Z., Klipfell, E., Bennett, B.J., Koeth, R., Levison, B.S., Dugar, B., Feldstein, A.E., Britt, E.B., Fu, X., Chung, Y.M., et al. (2011). Gut flora metabolism of phosphatidylcholine promotes cardiovascular disease. *Nature* **472**, 57–63.
17. Falls, J.G., Ryu, D.Y., Cao, Y., Levi, P.E., and Hodgson, E. (1997). Regulation of mouse liver flavin-containing monooxygenases 1 and 3 by sex steroids. *Arch. Biochem. Biophys.* **342**, 212–223.
18. Bain, M.A., Fornasini, G., and Evans, A.M. (2005). Trimethylamine: metabolic, pharmacokinetic and safety aspects. *Curr. Drug Metab.* **6**, 227–240.
19. al-Waiz, M., Mikov, M., Mitchell, S.C., and Smith, R.L. (1992). The exogenous origin of trimethylamine in the mouse. *Metabolism* **41**, 135–136.
20. Runck, A.M., Moriyama, H., and Storz, J.F. (2009). Evolution of duplicated beta-globin genes and the structural basis of hemoglobin isoform differentiation in Mus. *Mol. Biol. Evol.* **26**, 2521–2532.
21. Kobayakawa, K., Kobayakawa, R., Matsumoto, H., Oka, Y., Imai, T., Ikawa, M., Okabe, M., Ikeda, T., Itohara, S., Kikusui, T., et al. (2007). Innate versus learned odour processing in the mouse olfactory bulb. *Nature* **450**, 503–508.
22. Fendt, M., Endres, T., Lowry, C.A., Apfelbach, R., and McGregor, I.S. (2005). TMT-induced autonomic and behavioral changes and the neural basis of its processing. *Neurosci. Biobehav. Rev.* **29**, 1145–1156.
23. Fendt, M. (2006). Exposure to urine of canids and felids, but not of herbivores, induces defensive behavior in laboratory rats. *J. Chem. Ecol.* **32**, 2617–2627.
24. Johnson, M.A., Tsai, L., Roy, D.S., Valenzuela, D.H., Mosley, C., Magklara, A., Lomvardas, S., Liberles, S.D., and Barnea, G. (2012). Neurons expressing trace amine-associated receptors project to discrete glomeruli and constitute an olfactory subsystem. *Proc. Natl. Acad. Sci. USA* **109**, 13410–13415.
25. Krueger, S.K., and Williams, D.E. (2005). Mammalian flavin-containing monooxygenases: structure/function, genetic polymorphisms and role in drug metabolism. *Pharmacol. Ther.* **106**, 357–387.
26. Shirangi, T.R., Dufour, H.D., Williams, T.M., and Carroll, S.B. (2009). Rapid evolution of sex pheromone-producing enzyme expression in *Drosophila*. *PLoS Biol.* **7**, e1000168.
27. Waxman, D.J., and Holloway, M.G. (2009). Sex differences in the expression of hepatic drug metabolizing enzymes. *Mol. Pharmacol.* **76**, 215–228.
28. Clodfelter, K.H., Holloway, M.G., Hodor, P., Park, S.H., Ray, W.J., and Waxman, D.J. (2006). Sex-dependent liver gene expression is extensive and largely dependent upon signal transducer and activator of transcription 5b (STAT5b): STAT5b-dependent activation of male genes and repression of female genes revealed by microarray analysis. *Mol. Endocrinol.* **20**, 1333–1351.
29. Gower, D.B., Holland, K.T., Mallet, A.I., Rennie, P.J., and Watkins, W.J. (1994). Comparison of 16-androstene steroid concentrations in sterile apocrine sweat and axillary secretions: interconversions of 16-androstanes by the axillary microflora—a mechanism for axillary odour production in man? *J. Steroid Biochem. Mol. Biol.* **48**, 409–418.
30. Schellinck, H.M., West, A.M., and Brown, R.E. (1992). Rats can discriminate between the urine odors of genetically identical mice maintained on different diets. *Physiol. Behav.* **51**, 1079–1082.
31. Zomer, S., Dixon, S.J., Xu, Y., Jensen, S.P., Wang, H., Lanyon, C.V., O'Donnell, A.G., Clare, A.S., Gosling, L.M., Penn, D.J., and Brereton, R.G. (2009). Consensus multivariate methods in gas chromatography mass spectrometry and denaturing gradient gel electrophoresis: MHC-congenic and other strains of mice can be classified according to the profiles of volatiles and microflora in their scent-marks. *Analyst (Lond.)* **134**, 114–123.
32. Zala, S.M., Potts, W.K., and Penn, D.J. (2004). Scent-marking displays provide honest signals of health and infection. *Behav. Ecol.* **15**, 338–344.
33. Liberles, S.D., Horowitz, L.F., Kuang, D., Contos, J.J., Wilson, K.L., Siltberg-Liberles, J., Liberles, D.A., and Buck, L.B. (2009). Formyl peptide receptors are candidate chemosensory receptors in the vomeronasal organ. *Proc. Natl. Acad. Sci. USA* **106**, 9842–9847.
34. Rivière, S., Challet, L., Fluegge, D., Spehr, M., and Rodriguez, I. (2009). Formyl peptide receptor-like proteins are a novel family of vomeronasal chemosensors. *Nature* **459**, 574–577.
35. Tizzano, M., Gulbransen, B.D., Vandenbeuch, A., Clapp, T.R., Herman, J.P., Sibhatu, H.M., Churchill, M.E., Silver, W.L., Kinnamon, S.C., and Finger, T.E. (2010). Nasal chemosensory cells use bitter taste signaling to detect irritants and bacterial signals. *Proc. Natl. Acad. Sci. USA* **107**, 3210–3215.
36. Jemiolo, B., Alberts, J., Sochinski-Wiggins, S., and Novotny, M. (1985). Behavioural and endocrine responses of female mice to synthetic analogues of volatile compounds in male urine. *Anim. Behav.* **33**, 1114–1118.
37. Lin, D.Y., Zhang, S.Z., Block, E., and Katz, L.C. (2005). Encoding social signals in the mouse main olfactory bulb. *Nature* **434**, 470–477.
38. Jemiolo, B., Xie, T.M., and Novotny, M. (1991). Socio-sexual olfactory preference in female mice: attractiveness of synthetic chemosignals. *Physiol. Behav.* **50**, 1119–1122.
39. Roberts, S.A., Simpson, D.M., Armstrong, S.D., Davidson, A.J., Robertson, D.H., McLean, L., Beynon, R.J., and Hurst, J.L. (2010). Darcin: a male pheromone that stimulates female memory and sexual attraction to an individual male's odour. *BMC Biol.* **8**, 75.
40. Del Punta, K., Leinders-Zufall, T., Rodriguez, I., Jukam, D., Wysocki, C.J., Ogawa, S., Zufall, F., and Mombaerts, P. (2002). Deficient pheromone responses in mice lacking a cluster of vomeronasal receptor genes. *Nature* **419**, 70–74.
41. Keller, A., Zhuang, H., Chi, Q., Vosshall, L.B., and Matsunami, H. (2007). Genetic variation in a human odorant receptor alters odour perception. *Nature* **449**, 468–472.